

BIOMATIK

Tel: (519) 489-7195, (800) 836-8089 Fax: (519) 231-0140, (877) 221-3515 Email: info@biomatik.com http://www.biomatik.com

CAU26858

Polyclonal Antibody to Cystatin C (Cys-C)
Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Traits: Liquid

Concentration: 0.36mg/ml

UOM: 200µl

Cross Reactivity: Mouse; Rat; Pig **Applications:** WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant Cys-C (Ser27~Ala146) expressed in E.coli

[APPLICATIONS]

Western blotting: 0.5-2µg/mL; Immunohistochemistry: 5-20µg/mL; Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]


Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]









Figure. Western BlotSample: Lane1: Human Serum; Lane2: Rat Brain Tissue.



DAB staining on IHC-P; Samples: Human Intestine Tissue; Primary Ab: 20µg/ml Rabbit Anti-Human CST3 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.